

Change Notification for the UK Blood Transfusion Services

No. 29 - 2023

Chronic Fatigue Syndrome

Revised entry

Changes are indicated using the key below. This formatting will not appear in the final entry.

original text

«inserted text»

The following changes apply to:

BM-DSG CB-DSG GDRI TD-DSG WB-DSG

Bone Marrow
& Peripheral
Blood Stem Cell Cord Blood

Cor

Also Known As	CFS, myalgic encephalomyelitis (ME), post-viral fatigue syndrome and Systemic Exertion Intolerance Disease (SEID).
Obligatory	Must not donate
«Discretionary	If donor gives a history of fatigue following a viral infection, e.g. Glandular fever, with no relapse of symptoms and it is at least six months since all symptoms resolved, accept.»
Additional Information	CFS is generally diagnosed by excluding other conditions and may follow an infection that may or may not have been viral and which may be carried by the affected individual.
	It is most common between the ages of 25 and 45 years and women are affected more often than men. It is associated with easily induced and prolonged episodes of fatigue often accompanied by other symptoms.
	The condition is relapsing by nature and donation may make symptoms worse, or provoke a relapse in an affected individual.
	«Post viral fatigue can occur after an acute viral infection. Symptoms of fatigue can last weeks or months and may follow a relapsing course. It is important that individuals have fully recovered before being accepted to donate.»
Reason for Change	«A discretion to accept donors who have fully recovered from fatigue associated with an acute viral infection has been added.»
	The proposed new name for this syndrome has been included.

Jego 25

Dr Stephen Thomas
Professional Director - JPAC

jpac@nhsbt.nhs.uk