







Change Notification UK National Blood Services No. 54 - 2020

Tuberculosis

These changes apply to the Whole Blood and Components Donor Selection Guidelines

Please make the following changes to the "1. Affected Individual" section of this entry:

1. Affected Individual

Obligatory Must not donate if

a) Infected.

b) Less than 24 months from confirmation of cure.

c) Under follow-up.

Discretionary Donors with a diagnosis of Latent TB can donate, as long as they are not

currently undergoing investigation or treatment.

Donors on antibiotic treatment for Latent TB only can donate 7 days after

their last dose.

See if Relevant For BCG immunization:

Immunization - Live
Tuberculin PPD Test

Latent TB

Additional Information Tuberculosis can be present in many tissues and be spread through the

blood stream. It is sensible to exclude people who may have active disease from donating to prevent any possibility of transmitting the infection.

non donating to provent any possibility of transmitting the infection.

Someone with Latent TB does not have active disease and can usually be accepted to donate. However, once diagnosed they may be offered a course of antibiotics against TB. This is to reduce the risk of future reactivation of their infection. Treatment is for either three months or six months. Donors with Latent TB should not be accepted while on treatment, but can return to

donate seven days after completion of antibiotics.

Page 1 of 2

\Continued

Tel: 0114 358 4903

Fax: 0114 358 4494









Tel: 0114 358 4903

Fax: 0114 358 4494

Email: caroline.smith@nhsbt.nhs.uk

Reason for change

Advice and background information on Latent TB has been added.

Changes to Keyword/Search:

- Direct 'Latent Tuberculosis' to the 'Tuberculosis' entry rather than 'Accept'
- 'Accept Entry': No change required to text, but remove 'Latent Tuberculosis' from keywords.

Smaclena

Dr Sheila MacLennan

Professional Director - Joint UKBTS Professional Advisory Committee